arXiv:1402.1875v2 [math.AT] 18 Apr 2014

THE LUSTERNIK-SCHNIRELMANN CATEGORY OF METRIC SPACES

TULSI SRINIVASAN

Abstract. We extend the theory of the Lusternik-Schnirelmann category to general metric spaces by means of covers by arbitrary subsets. We also generalize the definition of the strict category weight. We show that if the Bockstein homomorphism on a metric space X is non-zero, then $\operatorname{cat} X \geq 2$, and use this to compute the category of Pontryagin surfaces. Additionally, we prove that a Polish space with LS-category n can be presented as the inverse limit of ANR spaces of category at most n.

1. Introduction

The *Lusternik-Schnirelmann category* (LS-category) of a space X is the smallest integer n such that X admits a cover $U_0, \ldots, U_n$ by open sets that are contractible in X. It is known that if the sets U_i in the definition are taken to be closed, then the category remains the same for ANR spaces. For general spaces, the definition of the LS-category by means of closed or open sets is not satisfactory. For instance, the category of the Menger curve is infinite if we consider covers of open or closed subsets. In [Sr] we considered arbitrary subsets of X in the definition of the LS-category, showed that the new definition coincides with the classical definition on compact metric ANRs, and applied it to fractal Peano continua like Menger spaces and Pontryagin surfaces. In this paper we discuss the LS-category of general metric spaces. In particular, we show that the definition with general subsets agrees with the classical one on all metric ANR spaces. Thus, we may use the notation $\operatorname{cat} X$ for the LS-category defined by means of arbitrary subsets. We note here that our general sets can be assumed to be G_δ on all complete metric spaces.

We then apply the general subsets definition to the category of maps. We recall that the *Lusternik-Schnirelmann category* (LS-category), $\operatorname{cat} f$, of a map $f : X \to Y$ is the smallest integer n such that $X = \bigcup_{i=0}^{n} U_i$,

Date: April 21, 2014.

2000 *Mathematics Subject Classification.* Primary 55M30; Secondary 57N65, 54F45.

where the U_i are open in X and each restriction $f|_{U_i}$ is nullhomotopic. Then the LS-category of a space X can be defined to be $\operatorname{cat} \mathrm{id}_X$. Also, if $A \subset X$, then the relative category of A is $\operatorname{cat}_X A = \operatorname{cat} \iota_A$, where $\iota : A \hookrightarrow X$ is the inclusion map. We generalize this definition by replacing open sets by arbitrary ones (see Definition 3.1), and study its properties. We use this to extend the theory of category weight to general metric spaces. We also prove an analogue to a generalization of Freudenthal's theorem in dimension theory, namely that a Polish metric space with LS-category n can be presented as the inverse limit of ANRs of category at most n.

Additionally, we apply the theory to Bockstein elements to give a lower bound for the LS-category of Pontryagin surfaces Π_p, which leads to the equality $\operatorname{cat} \Pi_p = 2$ for all p. Note that the case $p = 2$ was dealt with in [Sr], by using cup-length as a lower bound.

2. Preliminaries

Recall that given a weakly hereditary topological class of spaces $\mathcal{C}$, a space $X \in \mathcal{C}$ is called an *absolute neighborhood retract* (ANR) *for* $\mathcal{C}$ if every homeomorphic image of X as a closed subset of any $Z \in \mathcal{C}$ is a neighborhood retract of Z. If X is an ANR for the class of metrizable spaces, then we will call X a metric/metrizable ANR.

The following result is known.

2.1. **Proposition.** *Let X be a metric space, A a subset of X, and $\mathcal{V}' = \{V_i'\}_{i\in\mathbb{N}}$ a cover of A by sets open in A with $diam(V_i') < \delta_i$. Then $\mathcal{V}'$ can be extended to a cover $\mathcal{V} = \{V_i\}_{i\in N}$ of A by sets open in X with the same nerve and such that, for all $i \in \mathbb{N}$, $V_i \cap A = V_i'$ and $diam(V_i) < 2\delta_i$.*

Proof. Set

$$V_i = \bigcup_{a\in V_i'} B(a, \frac{1}{2}\min\{\delta_i, d(a, A - V_i')\}).$$

□

2.2. **Theorem** (Eilenberg-Kuratowski-Wojdyslawski Theorem)**.** [B, Hu] *Every bounded metric space can be embedded as a closed subset of a convex subset of a Banach space.*

The following is a version of a lemma that appears in [Wa].

2.3. **Theorem** (Modified Walsh Lemma)**.** *Let X be a metric space, A a subset of X, K a metrizable ANR, and $f : A \to K$ a map. Then, for any $\epsilon : K \to (0, \infty)$ (not necessarily continuous), there is an open set $U \supset A$ and a map $g : U \to K$ such that:*

(1) *For every $u \in U$, there exists $a_u \in A$ such that $d(g(u), f(a_u)) \leq \epsilon(f(a_u))$*
(2) *$g|_A$ is homotopic to f.*

Proof. By Theorem 2.2 (replacing the metric by an equivalent bounded metric if necessary), we can embed K as the closed subset of a convex subset T of a Banach space Y. Since K is an ANR, there is an open neighborhood O of K in T, and a retraction $r : O \to K$. For $y \in K$, we use the notation $B_T(y, \eta)$ to denote the convex open subset $B(y, \eta) \cap T$ of T.

For every $y \in K$, pick $\delta_y > 0$ such that:

(i) $B_T(y, 2\delta_y) \subset O$,
(ii) For all $y_1, y_2 \in B_T(y, 2\delta_y)$, we have $d(r(y_1), r(y_2)) < \epsilon(y)$.

For each $a \in A$, pick a neighborhood U_a of a that is open in A so that $f(U_a) \subset B_T(f(a), \delta_{f(a)})$. Let $\mathcal{V}' = \{V_i'\}$ be a locally finite refinement of the collection of U_a, and $\mathcal{V} = \{V_i\}$ the cover obtained by applying Proposition 2.1 to $\mathcal{V}'$. Let $U = \bigcup_i V_i$. Note here that we can take the U_a to have as small a diameter as necessary, so the diameter of the V_i can be made as small as required. This will be made use of in Lemma 2.5.

For each i, fix $a_i \in V_i'$ such that $V_i' \subset U_{a_i}$. Let $\{f_i : V_i \in \mathcal{V}\}$ be a partition of unity dominated by $\mathcal{V}$. Define $h : U \to Y$ by $h(u) = \sum_i f_i(u) f(a_i)$.

Choose any $u \in U$. Then u lies in precisely k elements of $\mathcal{V}$, say in $V_{i_1}, ..., V_{i_k}$. Assume that $\delta_{f(a_{i_j})} = \max\{\delta_{f(a_{i_1})}, ..., \delta_{f(a_{i_k})}\}$. Then Proposition 2.1 ensures that $f(a_{i_l}) \in B_T(f(a_{i_j}), 2\delta_{f(a_{i_j})})$ for all l, so $h(u)$ lies in this convex set as well. This means that h is a map from U to O.

Define $g : U \to K$ by $g = r \circ h$. If $u \in U$, then we have seen that $h(u)$ lies in $B_T(f(a_u), 2\delta_{f(a_u)})$ for some $a_u \in A$, so $d(g(u), f(a_u)) < \epsilon(f(a_u))$.

For every $a \in A$, $h(a)$ and $f(a)$ lie in some convex set, so $h|_A$ is homotopic to f in O via the straight line homotopy. The composition of this homotopy with r is then a homotopy in K between $g|_A$ and f. □

If Y is a metric ANR with a given open cover $\mathcal{U} = \{U_\lambda : \lambda \in \Lambda\}$, then there exists a refinement $\mathcal{V}$ of $\mathcal{U}$ such that given any X and any $f, g : X \to Y$ such that f and g are $\mathcal{V}$-close, the maps f and g are $\mathcal{U}$-homotopic [Hu]. We require a far less general version of this fact:

2.4. **Theorem.** [Hu] *Let K be a metric ANR. Then there exists a (not necessarily continuous) map $\epsilon : K \to (0, \infty)$ such that the covering $\mathcal{W} = \{B(y, \epsilon(y)) : y \in K\}$ has the following property: for any metric space X, any $\mathcal{W}$-close maps $f, g : X \to K$ are homotopic to each other.*

Let K be a metric ANR, and PK the path space $\{\phi : [0,1] \to K | \phi(1) = y_0\}$ for some $y_0 \in K$, equipped with the sup norm. For any $\epsilon : K \to (0,\infty)$, define $\bar{\epsilon} : PK \to (0,\infty)$ by $\bar{\epsilon}(\phi) = \epsilon(\phi(0))$. A metrizable space is an ANR for metrizable spaces iff it is an absolute neighborhood extensor (ANE) for metrizable spaces [Hu], from which it follows quite easily that if K is a metric ANR, then so is PK.

2.5. **Lemma.** *Let X be a metric space, $A \subset X$, K a metric ANR, and $f : X \to K$ a map such that the restriction of f to A is nullhomotopic. Then there exists $U \supset A$ open in X such that the restriction of f to U is also nullhomotopic.*

Proof. Let D be the metric on K and d the metric on X. Let PK be the path space $\{\phi : [0,1] \to K | \phi(1) = y_0\}$, under the sup metric D'. Let $\epsilon : K \to (0,\infty)$ be a map as in Theorem 2.4 and $\bar{\epsilon} : PK \to (0,\infty)$ as defined above.

Since $f|_A$ is nullhomotopic, there is a map $F : A \to PK$ satisfying $F(a)(0) = f(a)$ for all $a \in A$. For each $a \in A$, pick $\delta_a > 0$ so that for all $x \in X$ with $d(x,a) < \delta_a$, we have $D(f(x), f(a)) < \epsilon(f(a))$. We apply Theorem 2.3 to F, and construct an open neighborhood U of A and a map $G : U \to PK$ such that for every $u \in U$, there exists $a_u \in A$ such that $D'(G(u), F(a_u)) < \bar{\epsilon}(F(a_u))$. As noted in the proof of Theorem 2.3, we can construct U so that $\operatorname{diam}(V_i) < \delta_{a_i}$ for all i, the a_i and V_i being as in the proof of Theorem 2.3, and a_u being one of the a_i.

Let $g : U \to K$ be given by $g(u) = G(u)(0)$. Then $D(g(u), f(a_u)) \leq \bar{\epsilon}(F(a_u)) = \epsilon(f(a_u))$ and $D(f(u), f(a_u)) < \epsilon(f(a_u))$, and so both $g(u)$ and $f(u)$ lie in $B(f(a_u), \epsilon(f(a_u)))$. It follows that $f|_U$ is homotopic to g, which, by definition, is nullhomotopic. □

3. Generalized category of maps

3.1. **Definition.** For a map $f : X \to Y$, define the *generalized LS-category* of f, $\operatorname{cat}_g f$, to be the smallest integer k such that $X = \bigcup_{i=0}^{k} A_i$, where each restriction $f|_{A_i}$ is nullhomotopic. If no such k exists, we say that the category is infinite.

By definition, we have $\operatorname{cat}_g X = \operatorname{cat}_g \operatorname{id}_X$.

3.2. **Theorem.** *If $f : X \to Y$ be a map between a metric space X and a metric ANR Y, then $\operatorname{cat}_g f = \operatorname{cat} f$. In particular, if X is a metric ANR, then $\operatorname{cat}_g X = \operatorname{cat} X$.*

Proof. Clearly $\operatorname{cat}_g f \leq \operatorname{cat} f$, and equality holds if $\operatorname{cat}_g f$ is infinite. Suppose $\operatorname{cat}_g f = n$. Then we can write $X = \bigcup_{i=0}^{n} A_i$, where each $f|_{A_i}$

is nullhomotopic. By Lemma 2.5, there exist open sets U_i containing A_i such that each $f|_{U_i}$ is nullhomotopic. It follows that $\operatorname{cat} f \leq n$. $\square$

In the light of this, we will use the notation $\operatorname{cat} X$ for $\operatorname{cat}_g X$ for the remainder of this paper. However, we continue to use the subscript when referring to the LS-category of maps. We now list some basic properties of cat_g. The proofs are omitted, as they are nearly identical to those for cat (see, for instance, [CLOT]).

3.3. **Proposition** (Properties of cat_g)**.** *i) For any map $f : X \to Y$, we have* $\operatorname{cat}_g f \leq \min\{\operatorname{cat} X, \operatorname{cat} Y\}$

ii) If $f : X \to Y$ and $h : Y \to Z$ are maps, then

$$\operatorname{cat}_g(h \circ f) \leq \min\{\operatorname{cat}_g f, \operatorname{cat}_g h\}$$

iii) If $f : X \to Y$ is a homotopy equivalence, then $\operatorname{cat}_g f = \operatorname{cat} X = \operatorname{cat} Y$

iv) If $F \xrightarrow{i} E \xrightarrow{p} B$ is a fibration, then

$$\operatorname{cat}(E) \leq (\operatorname{cat}_g(i) + 1)(\operatorname{cat}_g(p) + 1) - 1.$$

Given a commutative ring R, the R-cup-length, $\text{cup-length}_R X$, of a space X is the smallest integer k such that all cup-products of length $k + 1$ vanish in the cohomology ring $H^*(X; R)$. We show that cup-length is a lower bound for cat_g.

3.4. **Theorem.** [MS, Theorem I.4.1 and Theorem 2.2.8, Appendix 1] *Every CW complex has the homotopy type of a metric ANR.*

3.5. **Theorem.** *Let $f : X \to Y$ be a map between metric spaces. Then for any commutative ring R,* $\operatorname{cat}_g(f) \geq \text{cup-length}_R(Im(f^*))$. *In particular,* $\operatorname{cat} X \geq \text{cup-length}_R X$.

Proof. We consider X and Y with the Alexander-Spanier cohomology or Čech cohomology (which coincide on paracompact spaces [HDD]). Let $\operatorname{cat}_g(f) = n$, and write $X = \cup_{i=0}^{n} A_i$, where each restriction $f|_{A_i}$ is nullhomotopic. Suppose $\text{cup-length}_R(Im(f^*)) > n$. Then there exist $\alpha_i \in H^{k_i}(Y; R)$ such that $f^*(\alpha_0) \smile ... \smile f^*(\alpha_n) \neq 0$. By the representability of the Čech cohomology, there exist maps $g_i : Y \to K(R, k_i)$ such that $g^*(u) = \alpha_i$, for a certain $u \in H^{k_i}(K(R, k_i); R)$. By Theorem 3.4 we may assume that the $K(R; k_i)$ are metrizable. Then each map $g_i \circ f : X \to K_i$ is a map between a metric space and a metric ANR with $g_i \circ f|_{A_i}$ nullhomotopic, so Lemma 2.5 implies that there exist

open sets $U_i \supset A_i$ such that $g_i \circ f|_{U_i}$ is nullhomotopic. Let $j_i : U_i \to X$ be the inclusion, and consider the following exact sequence:

$$\begin{array}{ccccccc} \cdots \to H^{k_i}(X, U_i; R) & \xrightarrow{q_i^*} & H^{k_i}(X; R) & \xrightarrow{j_i^*} & H^{k_i}(U_i; R) & \longrightarrow & \cdots \\ & & f^* \uparrow & & & & \\ & & H^{k_i}(Y; R) & & & & \end{array}$$

We have $j_i^*(f^*(\alpha_i)) = (g_i \circ f \circ j)^*(u) = 0$, so exactness implies that there exist $\bar{\alpha}_i \in H^{k_i}(X, U_i; R)$ such that $q_i^*(\bar{\alpha}_i) = f^*(\alpha_i)$. The rest of the proof follows as in the proof of the analogous statement for cat (see [CLOT]). Namely, in view of the relative cup-product formula for the Alexander-Spanier cohomology (see [Sp], p. 315),

$$H^k(X, U; R) \times H^l(X, V; R) \xrightarrow{\cup} H^{k+l}(X, U \cup V; R),$$

and the fact that $H^n(X, X; R) = 0$, we obtain a contradiction.

□

We conclude this section with a result analogous to the following generalization of Freudenthal's theorem in dimension theory: an n-dimensional completely metrizable space can be expressed as the inverse limit of polyhedra of dimension at most n [Na].

3.6. **Theorem.** *Let X be a Polish space with $\operatorname{cat} X \leq n$. Then X can be presented as the inverse limit of a sequence of ANR spaces, each of which has category $\leq n$.*

Proof. Suppose $X = \cup_{i=0}^n A_i$, where each A_i is contractible in X. Embed X as a closed subset of $\mathbb{R}^\omega$ [Ch], and take an open neighborhood U of X. By Lemma 2.5, we can enlarge each A_i to $U_{1,i}$ that is open and contractible in U, with $U_{1,i} \subset N_1(A_i)$. Set $U_1 = \bigcup_i U_{1,i}$ and $L_1 = \bigcup_i CU_{1,i}$, where $CY = (Y \times [0,1])/(Y \times \{1\})$ denotes the cone on Y. Then U_1 and L_1 are ANR spaces with $\operatorname{cat} L_1 \leq n$.

Next, we expand each A_i to $U_{2,i}$ that is open and contractible in U_1, with $U_{2,i} \subset N_{1/2}(A_i)$. Set $U_2 = \bigcup_i U_{2,i}$ and $L_2 = \bigcup_i CU_{2,i}$. Then U_2 and L_2 are ANRs with $\operatorname{cat} L_2 \leq n$. Since each $U_{2,i}$ is contractible in U_1, there is a map $H_{2,i} : CU_{2,i} \to U_1$ with $H_{2,i}|_{U_{2,i} \times \{0\}} = \mathrm{id}_{U_{2,i}}$. We paste these together to obtain a map $q_2 : L_2 \to L_1$ with $q_2|_{U_2} = \mathrm{id}_{U_2}$. Continuing this way, we obtain the inverse system $\{L_j, q_j\}$ of ANR spaces of dimension $\leq n$, which we claim has inverse limit L identical to X.

Since X is complete and each $U_{n,i} \subset N_{1/n}(A_i)$, the inverse limit of the inverse sequence $\{U_j, q_j|_{U_j}\}$ is X. Thus the maps $q_n : L_n \to U_{n-1} \subset L_{n-1}$ induce a map $q : L \to X$. Also, the inclusions $X \hookrightarrow L_n$ induce

$p : X \to L$ with $q \circ p = \mathrm{id}_X$, which implies that q is surjective. Finally, suppose two elements (y_j) and (z_j) in L are both mapped by q to $(x_j) \in X$. Then for each n we have $y_n = q_{n+1}(y_{n+1}) = x_n = q_{n+1}(z_{n+1}) = z_n$, so q is injective as well.

□

4. Category and the Bockstein homomorphism

The Bockstein homomorphism on a space X is the map

$$\beta_X : H^1(X; \mathbb{Z}_p) \to H^2(X; \mathbb{Z}_p)$$

obtained from the exact sequence $0 \to \mathbb{Z}_p \xrightarrow{p} \mathbb{Z}_{p^2} \to \mathbb{Z}_p \to 0$. It is known that for a metric ANR X, if $\beta_X \neq 0$, then $\operatorname{cat} X \geq 2$. [FH, Ru]

For metric ANR spaces, the usual definition of category coincides with Ganea's definition, i.e., given $f : X \to Y$, $\operatorname{cat} f \leq n$ iff there exists a map $\tilde{f} : X \to G_n(Y)$ such that $p_n \circ \tilde{f} \simeq f$, where $G_n(Y)$ is the nth Ganea space of Y and $p_n : G_n(Y) \to Y$ the nth Ganea map [CLOT]. The Ganea definition of category, together with the fact that $G_n(K(\Pi, 1))$ has the homotopy type of a CW-complex of dimension $\leq n$ for any group Π, yields a proof that $\beta_X \neq 0 \Rightarrow \operatorname{cat} X \geq 2$. It is not clear whether or not the Ganea definition can be applied to general spaces, but we can modify the technique above to obtain the same lower bound on the LS-category of a general metric space.

4.1. **Theorem.** *Let X be a metrizable space on which the Bockstein homomorphism β_X is non-zero. Then* $\operatorname{cat} X \geq 2$.

Proof. Since $\beta_X : H^1(X; \mathbb{Z}_p) \to H^2(X; \mathbb{Z}_p) \neq 0$, there exists $\alpha \in H^1(X; \mathbb{Z}_p)$ such that $\beta_X(\alpha) \neq 0$. By the representability of the Čech cohomology, there is a map $f : X \to K(\mathbb{Z}_p, 1)$ such that $f^*(i) = \alpha$, for a certain $i \in H^1(K(\mathbb{Z}_p, 1); \mathbb{Z}_p)$.

By Theorem 2.2, we can embed X as a closed subset of a convex subset T of a Banach space. Since $K(\mathbb{Z}_p, 1)$ is an ANE, f extends to a map $\tilde{f} : L \to K(\mathbb{Z}_p, 1)$ over some open neighborhood L of X in T. Since T is a convex subset of a Banach space, the Dugundji extension theorem [Du] implies that it is an ANR (in fact, an AR), and therefore L is also an ANR [Hu].

Suppose, for a contradiction, that $\operatorname{cat} X < 2$. Then $X = A_1 \cup A_2$, where the A_i are contractible in X. By Lemma 2.5, the A_i can be expanded to open sets U_i that are contractible in L. Set $K = U_1 \cup U_2$. Let $\iota_K : K \to L$ and $\iota_X : X \to K$ be the inclusion maps, and let $\bar{f} : K \to K(\mathbb{Z}_p, 1)$ be the restriction of $\tilde{f}$ to K. We claim that the Bockstein homomorphism $\beta_K : H^1(K; \mathbb{Z}_p) \to H^2(K; \mathbb{Z}_p)$ is non-zero.

Let $\bar{\alpha} = \bar{f}^*(i)$. From the diagram below, we have $0 \neq \beta_X(\alpha) = \beta_X \circ f^*(i) = \beta_X \circ \iota_X^* \circ \bar{f}^*(i) = \iota_X^* \circ \beta_K \circ \bar{f}^*(i) = \iota_X^*(\beta_K(\bar{\alpha}))$.

$$\begin{array}{ccc} H^1(K;\mathbb{Z}_p) & \xrightarrow{\iota_X^*} & H^1(X;\mathbb{Z}_p) \\ \downarrow{\scriptstyle \beta_K} & & \downarrow{\scriptstyle \beta_X} \\ H^2(K;\mathbb{Z}_p) & \xrightarrow{\iota_X^*} & H^2(X;\mathbb{Z}_p) \end{array}$$

This proves the claim.

However, since $\operatorname{cat} \iota_K < 2$ and K, L are metric ANRs, there exists a section s from K to the Ganea space $G_1(L)$, which induces $s^* : H^2(G_1(L);\mathbb{Z}_p) \to H^2(K;\mathbb{Z}_p)$. Since $G_1(K(\mathbb{Z}_p,1))$ has the homotopy type of a cell-complex of dimension ≤ 1, $H^2(G_1(K(\mathbb{Z}_p,1));\mathbb{Z}_p) = 0$, so i is mapped to 0 in $H^2(K;\mathbb{Z}_p)$ under $s^* \circ G_1(\tilde{f}^*) \circ p_1^* \circ \beta$ (see the diagram below). This implies that $\beta_K(\bar{\alpha}) = 0$, which is a contradiction.

$$\begin{array}{ccccc} H^2(G_1(K(\mathbb{Z}_p,1));\mathbb{Z}_p) & \xrightarrow{G_1(\tilde{f}^*)} & H^2(G_1(L);\mathbb{Z}_p) & \xrightarrow{G_1(\iota_K^*)} & H^2(G_1(K);\mathbb{Z}_p) \\ \uparrow{\scriptstyle p_1^*} & & \uparrow{\scriptstyle p_{1,L}^*} & \overset{s^*}{\swarrow} & \uparrow{\scriptstyle p_{1,K}^*} \\ H^2(K(\mathbb{Z}_p,1);\mathbb{Z}_p) & \xrightarrow{\tilde{f}^*} & H^2(L;\mathbb{Z}_p) & \xrightarrow{\iota_K^*} & H^2(K;\mathbb{Z}_p) \\ \uparrow{\scriptstyle \beta} & & \uparrow{\scriptstyle \beta_L} & & \uparrow{\scriptstyle \beta_K} \\ H^1(K(\mathbb{Z}_p,1);\mathbb{Z}_p) & \xrightarrow{\tilde{f}^*} & H^1(L;\mathbb{Z}_p) & \xrightarrow{\iota_K^*} & H^1(K;\mathbb{Z}_p) \end{array}$$

□

We use this result to compute the category of the Pontryagin surface Π_p, which is constructed as follows: let M_p be the mapping cylinder of the degree p map $f_p : S^1 \to S^1$, and let $\partial M_p \subset M_p$ denote the domain of f_p. Let $L_1 = S^2$, and suppose L_{k-1} has been constructed for $k \geq 2$. To construct L_k, take a sufficiently small triangulation of L_{k-1}, remove the interior of each of the triangles Δ and identify each $\partial\Delta$ with a copy of ∂M_p. Define $q_{k+1} : L_{k+1} \to L_k$ to be the map that collapses each $image(f_p) \subset M_p$ to a single point. Π_p is defined to be the inverse limit of $\{L_k, q_{k+1}\}$.

4.2. **Theorem.** *The category of the Pontryagin surface* $\Pi_p = 2$.

Proof. We refine the construction above by attaching copies of M_p one at a time as follows: Let M_p be as above, and let $X_1 = L_1 = S_2$. To construct L_2, we triangulate S_2 with, say, n_1 triangles. Let X_2 be the space obtained when one of the triangles Δ has its interior removed and its boundary identified with ∂M_p. Define $\lambda_2 : X_2 \to X_1$ to be the map that collapses $image(f_p) \subset M_p$ to a point. Let X_3 be obtained

from X_2 by removing the interior of a second triangle and attaching a second copy of M_p, and so on. We get $X_{n_1} = L_2$, and then use the same process to construct X_{n_1+1}. Let $\lambda_{k+1} : X_{k+1} \to X_k$ be the map that collapses $image(f_p)$ in the copy of M_p added to X_k to form X_{k+1} to a single point. Then Π_p is the inverse limit of $\{X_k, \lambda_{k+1}\}$.

We claim that the Bockstein homomorphism on β on Π_p is non-zero. This, along with the fact that $\operatorname{cat} X \leq \dim X$ [Sr], will imply that $\operatorname{cat} \Pi_p = 2$.

The Mayer-Vietoris sequence can be used to show that the maps $\lambda^*_{k+1} : H^2(X_k; \mathbb{Z}_p) \to H^2(X_{k+1}; \mathbb{Z}_p)$ are isomorphisms on $\mathbb{Z}_p$ for all k [Dr], and the long exact sequence for cohomology implies that each $\lambda^*_{k+1} : H^1(X_k; \mathbb{Z}_p) \to H^1(X_{k+1}; \mathbb{Z}_p)$ is injective. Also, since $H^2(X_2)$ has one $\mathbb{Z}_p$ component, the Bockstein map $\beta_2 : H^1(X_2; \mathbb{Z}_p) \to H^2(X_2; \mathbb{Z}_p)$ is an isomorphism [H], so the commutative diagram below can be used to show, inductively, that each $\beta_k : H^1(X_k; \mathbb{Z}_p) \to H^2(X_k; \mathbb{Z}_p)$ is non-zero:

$$\begin{array}{ccc} H^1(X_n; \mathbb{Z}_p) & \xrightarrow{\beta_n} & H^2(X_n; \mathbb{Z}_p) \\ \downarrow{\scriptstyle \lambda^*_{n+1}} & & \downarrow{\scriptstyle \cong} \\ H^1(X_{n+1}; \mathbb{Z}_p) & \xrightarrow{\beta_{n+1}} & H^2(X_{n+1}; \mathbb{Z}_p) \end{array}$$

□

5. Category Weight

For a metric space X the *strict category weight*, $\operatorname{swgt}(u)$, of $0 \neq u \in H^*(X; R)$, is the largest integer k such that for every map $\phi : A \to X$, where A is a metric space and $\operatorname{cat} \phi < k$, we have $\phi^*(u) = 0$. The strict category weight was first introduced in [Ru], in which X and A are assumed to be CW complexes, and in which it is shown that $\operatorname{cat} X \geq \operatorname{swgt}(u)$ for every $0 \neq u \in H^*(X; R)$. Strict category weight is a homotopy invariant version of Fadell and Husseini's category weight [FH], which is also a lower bound for the LS-category of a space. The definition can also be given with X and A paracompact Hausdorff spaces, as in [Ru1]. We generalize this as follows:

5.1. **Definition.** Let X be a metric space and let $0 \neq u \in H^*(X; R)$. Define the *generalized strict category weight of* u, $\operatorname{swgt}_g(u)$, *by*

$$\operatorname{swgt}_g(u) \geq k \Leftrightarrow \phi^*(u) = 0 \text{ for all } \phi : A \to X \text{ with } A \text{ metric, } \operatorname{cat}_g \phi < k.$$

Here R is a commutative ring, and $H^*(X; R)$ is the Alexander-Spanier or Čěch cohomology.

5.2. **Proposition.** *Let X be a metric space and $0 \neq u \in H^*(X;R)$. Then*

i) If $f : Y \to X$ is such that $f^*(u) \neq 0$, then $\operatorname{swgt}_g(u) \leq \operatorname{cat}_g f$. In particular, $\operatorname{swgt}_g(u) \leq \operatorname{cat} X$.
ii) If X is an ANR, then $\operatorname{swgt}_g(u) = \operatorname{swgt}(u)$.

Proof. i) This follows from definition.
ii) Clearly, $\operatorname{swgt}_g(u) \leq \operatorname{swgt}(u)$. Suppose X is an ANR and $\operatorname{swgt}(u) = k$. Let $\phi : A \to X$ be a map from a metric space A with $\operatorname{cat}_g \phi < k$. By Lemma 2.5, $\operatorname{cat} \phi < k$, and hence $\phi^*(u) = 0$. □

5.3. **Proposition.** *Let $u, v \in H^*(X;R)$. Then*

$$\operatorname{swgt}_g(u \smile v) \geq \operatorname{swgt}_g(u) + \operatorname{swgt}_g(v).$$

Proof. Let $\operatorname{swgt}_g(u) = m$ and $\operatorname{swgt}_g(v) = n$, and consider a map $\phi : A \to X$ with $\operatorname{cat}_g \phi < m + n$. We need to show that $\phi^*(u \smile v) = 0$.

Let $u = f^*(u')$ and $v = g^*(v')$ for $f : X \to K(R, k_1)$ and $g : X \to K(R, k_2)$, and suppose $A = B_1 \cup ... \cup B_n \cup C_1 \cup ... \cup C_m$, where the restriction of ϕ to each B_i and C_j is nullhomotopic. Let $B = \bigcup_{i=1}^n B_i$ and $C = \bigcup_{j=1}^m C_j$. Since $\operatorname{cat}_g(\phi|_B) < n$, we have $(\phi|_B)^*(u) = 0$, and hence the restriction of $f \circ \phi$ to B is nullhomotopic. Then Lemma 2.5 gives us an open set $U \supset B$, the restriction of $f \circ \phi$ to which is nullhomotopic. Similarly, we obtain $V \supset C$ such that the restriction of $g \circ \phi$ to V is nullhomotopic.

Consider the exact sequence associated with (A, U):

$$\cdots \to H^{k_1}(A, U; R) \xrightarrow{q_1^*} H^{k_1}(A; R) \xrightarrow{j^*} H^{k_1}(U; R) \longrightarrow \cdots$$

Since $j^*\phi^*(u) = 0$, there exists a $\bar{u}$ with $q_1^*(\bar{u}) = \phi^*(u)$. Similarly, there exists a $\bar{v} \in H^{k_2}(A, V; R)$ with $q_2^*(\bar{v}) = \phi^*(v)$. Now $\bar{u} \smile \bar{v} = 0$, as it lies in $H^*(A, U \cup V; R)$. But $\bar{u} \smile \bar{v} \mapsto \phi^*(u) \smile \phi^*(v) = \phi^*(u \smile v)$, proving the claim.

□

5.4. **Proposition.** *Let X be a metric space. Then*
i) If $f : Y \to X$ is such that $f^*(u) \neq 0$ for some $u \in H^*(X;R)$, then $\operatorname{swgt}_g(f^*(u)) \geq \operatorname{swgt}_g(u)$. If f is a homotopy equivalence, then $\operatorname{swgt}_g(f^*(u)) = \operatorname{swgt}_g(u)$.
ii) For $0 \neq w \in H^s(K(\Pi, 1); R)$, we have $\operatorname{swgt}_g(w) \geq s$.
iii) If $f : X \to K(\Pi, 1)$ and $0 \neq f^*(w) = u \in H^s(X;R)$, then $\operatorname{swgt}_g(u) \geq s$.
iv) If $\beta_X(u) \neq 0$ for some $u \in H^1(X; \mathbb{Z}_p)$, then $\operatorname{swgt}_g(\beta_X(u)) \geq 2$.

Proof. i) Suppose $\mathrm{swgt}_g(u) = n$, and consider $\phi : A \to Y$ with $\mathrm{cat}_g\, \phi < n$. By Proposition 3.3, $\mathrm{cat}_g(f \circ \phi) < n$, so $0 = (f \circ \phi)^*(u) = \phi^*(f^*(u))$. The second statement follows from the first.

ii) By Theorem 3.4, we can pick a metric $K(\Pi, 1)$ space. Then $\mathrm{swgt}_g(u) = \mathrm{swgt}(u)$. It is known that if $0 \neq u \in H^s(K(\Pi, 1); R)$, then $\mathrm{swgt}(u) \geq s$ [CLOT].

iii) This follows from i) and ii).

iv) Suppose $u = f^*(i)$, where $f : X \to K(\mathbb{Z}_p, 1)$. Then $\beta_X(u) = f^*(\beta_{K(\mathbb{Z}_p,1)}(i))$, so we can apply iii). □

Note that iv), together with the inequality $\mathrm{swgt}_g(u) \leq \mathrm{cat}\, X$, provides an alternative proof of Theorem 4.1.

6. Acknowledgement

I am grateful to my advisor, Alexander Dranishnikov, for all his ideas and insights, and also to Yuli Rudyak for suggesting that I study category weight.

References

[B] K. Borsuk, *Theory of Retracts.* PWN, 1967.

[Ch] A. Chigogidze, *Inverse spectra.* North-Holland Publishing Co., 1996.

[CLOT] O. Cornea, G. Lupton, J. Oprea and D. Tanré, *Lusternik-Schnirelmann category.* AMS, 2003.

[Dr] A. N. Dranishnikov, Cohomological dimension theory of compact metric spaces. Topology Atlas Invited Contributions (6) no. 1, 2001.

[Du] J. Dugundji, *An extension of Tietze's theorem.* Pacific J. Math. 1, 1951.

[FH] E. Fadell and S. Husseini, *Category weight and Steenrod operations.* Papers in honor of José Adem (Spanish). Bol. Soc. Mat. Mexicana (2), 37, no. 1-2, 1992.

[H] A. Hatcher, *Algebraic Topology.* Cambridge University Press, 2002.

[Hu] S. T. Hu, *Theory of Retracts.* Wayne State Univ. Press, 1965.

[HDD] W. Hurewicz, J. Dugundji, and C. H. Dowker, *Continuous connectivity groups in terms of limit groups.* Ann. of Math. (2) 49, 1948.

[MS] S. Mardešić and J. Segal, *Shape theory. The inverse system approach.* North-Holland Publishing Co., 1982.

[Na] K. Nagami, *Dimension theory,* With an appendix by Yukihiro Kodama. Pure and Applied Mathematics, Vol. 37, Academic Press, 1970.

[Ru] Y. B. Rudyak, *On category weight and its applications.* Topology, 38, no. 1, 1999.

[Ru1] Y. B. Rudyak, *On strict category weight, gradient-like flows, and the Arnold conjecture.* Internat. Math. Res. Notices, no. 5, 2000.

[Sp] E. Spanier, *Algebraic Topology.* McGraw-Hill Book Co., 1966.

[Sr] T. Srinivasan *On the Lusternik-Schnirelmann category of Peano continua.* Topology and its Applications, 160, no. 13, 2013.

[Wa] J. J. Walsh, *Dimension, cohomological dimension, and cell-like mappings.* Lecture Notes in Math., 870, Springer, 1981.

Tulsi Srinivasan, Department of Mathematics, University of Florida, 358 Little Hall, Gainesville, FL 32611-8105, USA
E-mail address: tsrinivasan@ufl.edu